
Technical Procedure for Amido Black

1.0 Purpose – This is a blood print development procedure.

2.0 Scope – This procedure applies to items of evidence, porous and non-porous, that contain impressions in blood that require enhancing.

3.0 Definitions – N/A

4.0 Equipment, Materials and Reagents

4.1 Equipment and Materials

- Protective gloves and apron/coat
- Face shield visor and/or safety goggles
- Magnetic stir bar and magnetic retriever
- Two (2) 2000 mL glass beakers
- One (1) 1000 mL glass beaker
- Dark, shatter-proof container

4.2 Reagents

- 5-Sulfosalicyclic acid
- Naphthalene 12B, Naphthol Blue Black, or Amido Black
- Sodium carbonate
- Formic acid
- Acetic acid
- Kodak Photo Flo 200 Solution
- Purified water

5.0 Procedure

5.1 Preparation

5.1.1 Pour 500 mL of purified water into a clean 2000 mL glass beaker with a magnetic stir bar.

5.1.2 Continue to stir the solution while adding the following in order:

- 5-Sulfosalicyclic acid (20 g)
- Naphthalene 12B, Naphthol Blue Black, or Amido Black (3 g)
- Sodium carbonate (3 g)
- Formic acid (50 mL)
- Acetic acid (50 mL)
- Kodak Photo Flo 200 Solution (12.5 mL)

5.1.3 Stir until all crystals are dissolved.

5.1.4 Dilute mixture to approximately 1 liter with purified water.

5.1.5 Solution may be stored indefinitely as there is no expiration period.

5.2 Application

5.2.1 Amido Black may be applied to evidence either by using a spray bottle or by dipping the evidence. Completely cover the target area and allow to develop for approximately 3 to 5 minutes. Rinse item with tap water to remove any excess reagent after development.

5.3 Standards and Controls – The Forensic Scientist shall produce a self-made test print to be processed concurrently with items of evidence.

5.4 Calibration – N/A

5.5 Sampling – N/A

5.6 Calculations – N/A

5.7 Uncertainty of Measurement - N/A

6.0 Limitations – Amido Black may stain lighter background surfaces.

6.1 Amido Black will not detect the normal constituents of latent impressions; therefore, Amido Black must be used in sequence with other processing techniques. It is essential to develop any traditional latent impressions prior to treatment with Amido Black. Careful treatment with powders before Amido Black will not interfere with Amido Black Processing. Ninhydrin, if used, must precede the application of Amido Black.

6.2 Some impressions in blood on porous surfaces such as paper or cardboard may be improved by using Physical Developer after the application of Amido Black. Washing the item in a warm detergent solution after the Physical Developer treatment will reduce the Amido Black background staining. Use 1 mL of Tergitol 7 per liter of purified water at 50 °C, if the item of evidence will withstand this temperature. Wash the item until the background stain is reduced to acceptable levels.

6.3 Amido Black solution may be kept indefinitely in a dark, shatter proof container.

7.0 Safety – Amido Black shall be used only in a well-ventilated area such as a fume hood. The glacial acetic acid and formic acid in the solution may be corrosive. Eye protection shall be worn when preparing and transporting the reagent, and when processing items with the solution.

7.1 This solution may also be used at crime scenes provided that proper safety precautions are applied; however, only use in well ventilated areas or utilize a fan to remove the fumes produced.

8.0 References

Hamm, E. "Enhancement and Development of Blood Prints." *74th Annual Educational Conference International Association for Identification*. (June 22, 1989): 1-11.

Kent, T. ed. *Manual of Fingerprint Development Techniques: A Guide to the Selection and Use of Processing for the Development of Latent Fingerprints*. Police Scientific Development Branch, London (July 1992).

Lee, H.C. “Methods of Latent Print Development.” *Proceedings of the International Forensic Symposium on Latent Prints*. (July 1987): 15 – 24.

Lennard, C.J., and P. Margot. “A Sequencing of Reagents for the Improved Visualization of Latent Fingerprints.” *Proceedings of the International Forensic Symposium on Latent Prints*. (July 1987): 141-142.

Manual of Fingerprint Development Techniques. (January 1986): 2-8.

Manual of Fingerprint Development Techniques: A Guide to the Selection and Use of Processes for the Development of Latent Fingerprints. Scientific Research and Development Branch, London (1986).

Merchant, B., and C. Tague. “Developing Fingerprints in Blood: A Comparison of Several Chemical Techniques.” *Journal of Forensic Identification*. Vol. 57, 1: 76 – 57 (2007).

Norkus, P.M., and K. Noppinger. “Enhanced Latent Prints in Blood with a New Staining Technique.” *Proceedings of the International Forensic Symposium on Latent Prints*. (July 1987): 147.

Trozzi, T.A., R.L. Schwartz, and M.L. Hollars. *Processing Guide for Developing Latent Prints*. (2000): 1-64.

9.0 Records – N/A

10.0 Attachments – N/A

Revision History		
Effective Date	Version Number	Reason
09/17/2012	1	Original Document
10/31/2013	2	Added issuing authority to header
10/19/2015	3	Changed all references of Photo Flo 600 to Photo Flo 200 Header – updated issuing authority
03/30/2017	4	Header Update – Removed Digital reference.
01/19/2018	5	5.2.1 – Moved requirement for test print to “Standards and Control.” Updated to magnetic stir bar throughout.
02/01/2019	6	Changed number references to be numerals only instead of spelled out throughout document 5.1.2: changed to “Continue to stir the solution” Changed “distilled” to “purified” throughout document